

Functional renormalization group approach to the Anderson impurity model

Lorenz Bartosch,¹ Hermann Freire,¹ Jose Juan Ramos Cardenas,^{1,2} and Peter Kopietz¹

¹*Institut für Theoretische Physik, Universität Frankfurt,
Max-von-Laue Strasse 1, 60438 Frankfurt, Germany*

²*Departamento Fisico-Matematico, Universidad Autonoma de San Luis Potosi,
Niño Artillero 140, 78290 San Luis Potosi, Mexico*

(Dated: November 17, 2008)

We develop a functional renormalization group approach which describes the low-energy single-particle properties of the Anderson impurity model up to intermediate on-site interactions $U \lesssim 15\Delta$, where Δ is the hybridization in the wide-band limit. Our method is based on a generalization of a method proposed by Schütz, Bartosch and Kopietz [Phys. Rev. B **72**, 035107 (2005)], using two independent Hubbard-Stratonovich fields associated with transverse and longitudinal spin fluctuations. Although we do not reproduce the exponentially small Kondo scale in the limit $U \rightarrow \infty$, the spin fluctuations included in our approach remove the unphysical Stoner instability predicted by mean-field theory for $U > \pi\Delta$. We discuss different decoupling schemes and show that a decoupling which manifestly respects the spin-rotational invariance of the problem gives rise to the lowest quasi-particle weight. To obtain a closed flow equation for the fermionic self-energy we also propose a new truncation scheme of the functional renormalization group flow equations using Dyson-Schwinger equations to express bosonic vertex functions in terms of fermionic ones.

PACS numbers: 72.15.Qm, 71.27.+a, 71.10.Pm

I. INTRODUCTION

The Anderson impurity model (AIM) is one of the most important model systems in condensed matter physics.¹ The model was originally proposed by Anderson² to describe the properties of local moments in metals. Its Hamiltonian describes a single correlated impurity which is coupled to a band of non-interacting conduction electrons,

$$\begin{aligned} \hat{H} = & \sum_{\mathbf{k}\sigma} (\epsilon_{\mathbf{k}} - \sigma h) \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} \\ & + \sum_{\sigma} (E_d - \sigma h) \hat{d}_{\sigma}^\dagger \hat{d}_{\sigma} + U \hat{d}_{\uparrow}^\dagger \hat{d}_{\uparrow} \hat{d}_{\downarrow}^\dagger \hat{d}_{\downarrow} \\ & + \sum_{\mathbf{k}\sigma} (V_{\mathbf{k}}^* \hat{d}_{\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} + V_{\mathbf{k}} \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{d}_{\sigma}). \end{aligned} \quad (1.1)$$

Here, $\hat{c}_{\mathbf{k}\sigma}$ annihilates a non-interacting conduction electron with momentum $\mathbf{k}$, energy dispersion $\epsilon_{\mathbf{k}}$ and spin projection σ , while the operator $\hat{d}_{\sigma}$ annihilates a localized correlated d -electron with atomic energy E_d and on-site repulsion U . The hybridization between the d -electrons and the conduction electrons is characterized by the hybridization energy $V_{\mathbf{k}}$. We have also included in Eq. (1.1) the Zeemann energy h associated with an external magnetic field.

The thermodynamic and spectral properties of the AIM can be calculated accurately by means of Wilson's numerical renormalization group^{3,4,5} (NRG), see Ref. [6] for a recent review. However, the calculation of the spectral function $A(\omega)$ of the d -electrons by means of the NRG requires some computational effort, in particular if one needs accurate results for arbitrary frequencies ω . Since the quantitative knowledge of the spectral function of the AIM for all ω is essential in the context of the dy-

namical mean-field theory describing strong correlations in realistic three-dimensional fermion systems,⁷ it is important to develop approximate analytical methods for calculating the spectral function of the AIM. Although several analytical approaches have been proposed to describe the strong coupling regime,^{8,9,10,11,12,13,14} a satisfactory analytical alternative which can compete with the NRG in the strong coupling regime has not been found. This has motivated us to develop a new functional renormalization group (FRG) approach to the AIM which is not based on the weak coupling truncation used by other authors.^{15,16} Our approach extends the collective field FRG developed in Refs. [17,18,19], which is based on the partial bosonization of the two-body interaction using suitable Hubbard-Stratonovich fields. For simplicity, we focus here on the local moment regime,¹ where the energy E_d of the d -level is located below the Fermi energy, but its double occupancy is prohibited by a strong on-site interaction U . In the limit $U \rightarrow \infty$ the low-energy properties of the AIM are in this regime identical to those of the Kondo model describing only the spin degree of freedom of the impurity. It is then natural to decouple the interaction of the AIM in terms of Hubbard-Stratonovich fields representing collective spin-fluctuations, which should be treated non-perturbatively to describe the strong-coupling regime.

Since we are interested in the correlation functions of the d -electrons, we simply integrate out the conduction electrons using the coherent state functional integral. The ratio of the partition functions with and without interaction at constant chemical potential μ and inverse temperature β can then be written as

$$\frac{\mathcal{Z}}{\mathcal{Z}_0} = \frac{\int \mathcal{D}[d, \bar{d}] e^{-S_0[\bar{d}, d] - S_U[\bar{d}, d]}}{\int \mathcal{D}[d, \bar{d}] e^{-S_0[\bar{d}, d]}}, \quad (1.2)$$

where the Gaussian part is given by

$$\begin{aligned} S_0[\bar{d}, d] &= \int_0^\beta d\tau \sum_\sigma \bar{d}_\sigma(\tau) [\partial_\tau + E_d - \mu - \sigma h] d_\sigma(\tau) \\ &+ \int_0^\beta d\tau \int_0^\beta d\tau' \sum_\sigma \bar{d}_\sigma(\tau) \Delta^\sigma(\tau - \tau') d_\sigma(\tau') \\ &= - \int_\omega \sum_\sigma [i\omega - \xi_0^\sigma - \Delta^\sigma(i\omega)] \bar{d}_{\omega\sigma} d_{\omega\sigma}, \end{aligned} \quad (1.3)$$

with

$$\xi_0^\sigma = E_d - \mu - \sigma h, \quad (1.4)$$

and the interaction is

$$S_U[\bar{d}, d] = U \int_0^\beta d\tau \bar{d}_\uparrow(\tau) d_\uparrow(\tau) \bar{d}_\downarrow(\tau) d_\downarrow(\tau). \quad (1.5)$$

Here, $\int_\omega = \frac{1}{\beta} \sum_\omega$ denotes summation over fermionic Matsubara frequencies. Later we shall take the limit $\beta \rightarrow \infty$ where $\int_\omega = \int \frac{d\omega}{2\pi}$. The Fourier transform of the Grassmann fields $d_\sigma(\tau)$ in frequency space is defined by

$$d_\sigma(\tau) = \int_\omega e^{-i\omega\tau} d_{\omega\sigma}, \quad (1.6)$$

and the hybridization function is

$$\Delta^\sigma(\tau) = \int_\omega e^{-i\omega\tau} \Delta^\sigma(i\omega), \quad (1.7)$$

with Fourier components

$$\Delta^\sigma(i\omega) = \sum_{\mathbf{k}} \frac{|V_{\mathbf{k}}|^2}{i\omega - \epsilon_{\mathbf{k}} + \mu + \sigma h}. \quad (1.8)$$

The 0 + 1-dimensional quantum field theory defined in Eqs. (1.3)–(1.5) will be the starting point of our further calculations presented below.

Let us briefly outline the rest of this work: In Sec. II we decouple the interaction (1.5) in the spin-singlet particle-hole channel using a complex Hubbard-Stratonovich field which describes transverse spin-flip fluctuations. By treating the resulting field theory on the level of the Gaussian approximation, we obtain results for the dynamic structure factor and the self-energy in the ladder approximation. Of course, only for small U can we expect the ladder approximation to give reliable results. For larger couplings we find a ferromagnetic Stoner instability, consistent with the Hartree-Fock approximation but in contrast to well-established results. To go beyond the simple ladder approximation, we use in Sec. III a collective field FRG approach for our Bose-Fermi theory and implement a simple truncation to close the set of flow equations. Our inclusion of transverse spin fluctuations based on the FRG removes the unphysical Stoner instability and gives results which qualitatively describe

the correct physics. However, although the quasiparticle weight vanishes with increasing interaction as it should, it does not obey the well-known Kondo scaling. To improve on these results, we introduce in Sec. IV a new FRG approach with partial bosonization in both the transverse and the longitudinal channel. We discuss the ambiguities with the distribution of weight among the two channels and show that in the manifestly spin-rotationally invariant case we get the strongest suppression of the quasiparticle weight. Finally, in Sec. V we summarize our results and discuss some open questions.

II. LADDER APPROXIMATION IN THE SPIN-SINGLET PARTICLE-HOLE CHANNEL

A. Hartree-Fock approximation and Stoner instability

If we treat the interaction (1.5) within the self-consistent Hartree-Fock approximation, we obtain the renormalized excitation energy $\xi^\sigma = \xi_0^\sigma + \delta\xi^\sigma$, with

$$\delta\xi^\sigma = U \int_\omega G_0^{\bar{\sigma}}(i\omega), \quad (2.1)$$

where we have used the notation $\bar{\sigma} = -\sigma$ for the spin label. The self-consistent Hartree-Fock Green function is

$$G_0^\sigma(i\omega) = \frac{1}{i\omega - \xi_0^\sigma - \Delta^\sigma(i\omega)}. \quad (2.2)$$

The corresponding self-energy can be written as

$$\delta\xi^\sigma = \frac{U}{2} [n - \sigma m], \quad (2.3)$$

where $n = n_\uparrow + n_\downarrow$ is the average occupation and $m = n_\uparrow - n_\downarrow$ is the average magnetization of the impurity level. For a sufficiently strong interaction, Hartree-Fock theory predicts a finite value of m even for $h \rightarrow 0$. In the wide-band limit, where the hybridization function can be approximated by

$$\Delta^\sigma(i\omega) = -i\Delta^\sigma \text{sgn } \omega, \quad (2.4)$$

the Hartree-Fock self-consistency equation for the moment m can be evaluated analytically at zero temperature. For $h = 0$ (where $\Delta^\sigma = \Delta$ is independent of the spin projection) and in the particle-hole symmetric case (where $E_d - \mu = -U/2$ and $n = 1$) the result is

$$m = \int_\omega \sum_\sigma \sigma G_0^\sigma(i\omega) = \frac{2}{\pi} \arctan\left(\frac{Um}{2\Delta}\right). \quad (2.5)$$

This equation has a ferromagnetic solution ($m \neq 0$) if

$$u_0 \equiv \frac{U}{\pi\Delta} > 1. \quad (2.6)$$

This ferromagnetic Stoner-instability is an artefact of the Hartree-Fock approximation,¹ which should be removed once fluctuation corrections are properly taken into account.

B. Hubbard-Stratonovich transformation in the spin-singlet particle-hole channel

In the strong coupling limit we expect that transverse spin fluctuations should play an important role to remove the Stoner instability.¹¹ It is therefore natural to decouple the interaction in the spin-singlet particle-hole channel with the help of a bosonic Hubbard-Stratonovich field, describing transverse spin fluctuations. To this end, we use the antisymmetry of the Grassmann fields to write the integrand in the interaction functional (1.5) as

$$U\bar{d}_\uparrow d_\uparrow \bar{d}_\downarrow d_\downarrow = -U(\bar{d}_\uparrow d_\downarrow)(\bar{d}_\downarrow d_\uparrow) = -U\bar{s}(\tau)s(\tau), \quad (2.7)$$

where we have defined the composite spin-flip fields,

$$\bar{s}(\tau) = \bar{d}_\uparrow(\tau)d_\downarrow(\tau), \quad s(\tau) = \bar{d}_\downarrow(\tau)d_\uparrow(\tau). \quad (2.8)$$

Introducing the complex bosonic Hubbard-Stratonovich fields $\chi(\tau)$ and $\bar{\chi}(\tau)$ conjugate to $\bar{s}(\tau)$ and $s(\tau)$, the ratio (1.2) of the interacting and non-interacting partition functions can be written as a functional integral over a six-component superfield $\Phi = [d_\uparrow, \bar{d}_\uparrow, d_\downarrow, \bar{d}_\downarrow, \chi, \bar{\chi}]$,

$$\frac{\mathcal{Z}}{\mathcal{Z}_0} = \frac{\int \mathcal{D}[\Phi] e^{-S_0[\Phi] - S_1[\Phi]}}{\int \mathcal{D}[\Phi] e^{-S_0[\Phi]}}. \quad (2.9)$$

The Gaussian part of the bare action is

$$S_0[\Phi] = -\int_\omega \sum_\sigma [G_0^\sigma(i\omega)]^{-1} \bar{d}_{\omega\sigma} d_{\omega\sigma} + \int_{\bar{\omega}} U^{-1} \bar{\chi}_{\bar{\omega}} \chi_{\bar{\omega}}, \quad (2.10)$$

and the interaction part can be written as

$$S_1[\Phi] = \int_{\bar{\omega}} [\bar{s}_{\bar{\omega}} \chi_{\bar{\omega}} + s_{\bar{\omega}} \bar{\chi}_{\bar{\omega}}] - \int_\omega \sum_\sigma \delta\xi^\sigma \bar{d}_{\omega\sigma} d_{\omega\sigma}, \quad (2.11)$$

where

$$s_{\bar{\omega}} = \int_0^\beta d\tau e^{i\bar{\omega}\tau} s(\tau) = \int_\omega \bar{d}_{\omega\downarrow} d_{\omega+\bar{\omega},\uparrow}, \quad (2.12)$$

$$\bar{s}_{\bar{\omega}} = \int_0^\beta d\tau e^{-i\bar{\omega}\tau} \bar{s}(\tau) = \int_\omega \bar{d}_{\omega+\bar{\omega},\uparrow} d_{\omega\downarrow}. \quad (2.13)$$

Because $[G_0^\sigma(i\omega)]^{-1}$ in the Gaussian part (2.10) of the action is by definition the inverse Hartree-Fock Green function (2.2), the Hartree-Fock self-energy $\delta\xi^\sigma$ should be subtracted from the interaction in Eq. (2.11) as a counterterm. The boson-fermion interaction in Eq. (2.11) can be written as

$$\begin{aligned} \int_{\bar{\omega}} [\bar{s}_{\bar{\omega}} \chi_{\bar{\omega}} + s_{\bar{\omega}} \bar{\chi}_{\bar{\omega}}] = \\ \int_{\bar{\omega}} \int_\omega \left[\Gamma_0^{(\bar{d}_\uparrow d_\downarrow \chi)}(\omega + \bar{\omega}, \omega, \bar{\omega}) \bar{d}_{\omega+\bar{\omega},\uparrow} d_{\omega\downarrow} \chi_{\bar{\omega}} \right. \\ \left. + \Gamma_0^{(\bar{d}_\downarrow d_\uparrow \bar{\chi})}(\omega - \bar{\omega}, \omega, \bar{\omega}) \bar{d}_{\omega-\bar{\omega},\downarrow} d_{\omega\uparrow} \bar{\chi}_{\bar{\omega}} \right], \end{aligned} \quad (2.14)$$

with the bare spin-flip vertices given by

$$\Gamma_0^{(\bar{d}_\uparrow d_\downarrow \chi)}(\omega + \bar{\omega}, \omega, \bar{\omega}) = \Gamma_0^{(\bar{d}_\downarrow d_\uparrow \bar{\chi})}(\omega - \bar{\omega}, \omega, \bar{\omega}) = 1. \quad (2.15)$$

A graphical representation of these vertices is shown in Fig. 1.




FIG. 1: (Color online) Graphical representation of the bare boson-fermion vertices of the action $S_1[\Phi]$ in Eq. (2.11). Solid arrows pointing into the vertices represent d_σ , while outgoing solid arrows represent $\bar{d}_\sigma$. The small arrows indicate the spin projections. The wavy arrows represent the spin-flip field χ (pointing into the black dots) and $\bar{\chi}$ (pointing out of the black dots). The direction of the arrows associated with the wavy lines is chosen such that the incoming arrow adds spin to the vertex.

C. Gaussian propagator of transverse spin-fluctuations

If we integrate in Eq. (2.9) over the fermion fields, we obtain the effective action $S_{\text{eff}}[\bar{\chi}, \chi]$ of the spin-flip fields $\chi, \bar{\chi}$. Expanding $S_{\text{eff}}[\bar{\chi}, \chi]$ to quadratic order in the fields (Gaussian approximation), we obtain

$$S_{\text{eff}}[\bar{\chi}, \chi] \approx \int_{\bar{\omega}} [F_{\text{LA}}^\perp(i\bar{\omega})]^{-1} \bar{\chi}_{\bar{\omega}} \chi_{\bar{\omega}}. \quad (2.16)$$

The inverse spin-flip propagator in ladder approximation (LA) is given by

$$[F_{\text{LA}}^\perp(i\bar{\omega})]^{-1} = U^{-1} - \Pi_0^\perp(i\bar{\omega}), \quad (2.17)$$

where $\Pi_0^\perp(i\bar{\omega})$ is the non-interacting dynamic spin-flip susceptibility

$$\Pi_0^\perp(i\bar{\omega}) = -\int_\omega G_0^\uparrow(i\omega) G_0^\downarrow(i\omega - i\bar{\omega}). \quad (2.18)$$

In the wide-band limit, where $\Delta^\sigma(i\omega)$ is given by Eq. (2.4), the integration in Eq. (2.18) can be performed analytically at zero temperature. In this work we only need $\Pi_0^\perp(i\bar{\omega})$ in the absence of an external magnetic field and spontaneous ferromagnetism, where $\xi^\sigma = \xi$ and $\Delta^\sigma = \Delta$ are independent of the spin projection σ ,

$$\Pi_0^\perp(i\bar{\omega}) = \frac{\Delta}{\pi|\bar{\omega}|(|\bar{\omega}| + 2\Delta)} \ln \left[\frac{\xi^2 + (|\bar{\omega}| + \Delta)^2}{\xi^2 + \Delta^2} \right]. \quad (2.19)$$

If we assume in addition particle-hole symmetry so that $\xi = \xi_0 + \delta\xi = 0$, then Eq. (2.19) further simplifies to

$$\begin{aligned} \pi\Delta\Pi_0^\perp(i\bar{\omega}) &= \frac{\ln[1 + |\bar{\epsilon}|]}{|\bar{\epsilon}|(1 + |\bar{\epsilon}|/2)} \\ &= 1 - |\bar{\epsilon}| + \frac{5}{6}\bar{\epsilon}^2 + \mathcal{O}(|\bar{\epsilon}|^3), \end{aligned} \quad (2.20)$$

where $\bar{\epsilon} = \bar{\omega}/\Delta$. Within the LA the spectral density $S_{\text{LA}}^\perp(\omega)$ of transverse spin-fluctuations (the dynamic structure factor) is then given by

$$\Pi_{\text{LA}}^\perp(i\bar{\omega}) = \frac{\Pi_0^\perp(i\bar{\omega})}{1 - U\Pi_0^\perp(i\bar{\omega})} = \int_0^\infty \frac{d\omega}{\pi} S_{\text{LA}}^\perp(\omega) \frac{2\omega}{\omega^2 + \bar{\omega}^2}, \quad (2.21)$$




FIG. 2: (Color online) Graph of the transverse spin structure factor $S_{\text{LA}}^{\perp}(\omega)$ in ladder approximation for $h = 0$, $u_0 = U/(\pi\Delta) = 0.9$ and particle-hole symmetry. The characteristic energy scale of transverse spin fluctuations is in this approximation given by $\omega_* \approx \Delta(1 - u_0)$.

or, equivalently,

$$\text{Im } \Pi_{\text{LA}}^{\perp}(\omega + i0) = \text{sgn } \omega S_{\text{LA}}^{\perp}(|\omega|). \quad (2.22)$$

A graph of $S_{\text{LA}}^{\perp}(\omega)$ for $u_0 = U/(\pi\Delta) = 0.9$ is shown in Fig. 2. In the regime $0 < 1 - u_0 \ll 1$ the LA predicts a well defined peak in the dynamic structure factor at the energy scale

$$\omega_* = \Delta(1 - u_0). \quad (2.23)$$

The width of the peak is of the order of Δ . In fact, in this regime the low-frequency behavior of $S_{\text{LA}}^{\perp}(\omega)$ can easily be obtained analytically. For $|\omega| \ll \Delta$ we find

$$\Pi_{\text{LA}}^{\perp}(i\bar{\omega}) = \frac{1}{\pi} \frac{1}{\omega_* + u_0|\bar{\omega}|}, \quad (2.24)$$

so that the corresponding structure factor is

$$S_{\text{LA}}^{\perp}(\omega) \approx \frac{1}{\pi} \frac{\omega}{\omega_*^2 + \omega^2}. \quad (2.25)$$

For $u_0 \rightarrow 1$ the energy scale ω_* vanishes, indicating an instability towards spontaneous ferromagnetism, as suggested by the Hartree-Fock approximation discussed in Sec. II A. Because we know that the AIM does not exhibit spontaneous ferromagnetism for arbitrary values of U , we expect that the LA is only accurate for $u_0 \ll 1$. Before employing more sophisticated FRG methods to include fluctuation corrections which remove this unphysical Stoner instability, it is instructive to consider the d -electron self-energy within the LA, which we shall do in the following subsection.

D. Fermionic self-energy in ladder approximation

The LA in the particle-hole channel gives the following self-energy for the d -electrons,

$$\Sigma^{\sigma}(i\omega) = \int_{\bar{\omega}} F_{\text{LA}}^{\perp}(i\bar{\omega}) G_0^{\bar{\sigma}}(i\omega - i\sigma\bar{\omega}). \quad (2.26)$$




FIG. 3: Diagrams contributing to the fermionic self-energy $\Sigma(i\omega)$ in particle-hole ladder approximation, see Eq. (2.26). Solid arrows denote fermionic Hartree-Fock Green functions and wavy lines denote the bare interaction. The first-order (Hartree) diagram is subtracted in Eq. (2.27).




FIG. 4: (Color online) Graph of the quasiparticle residue Z as a function of $u_0 = U/(\pi\Delta)$ within LA for $h = 0$ and particle-hole symmetric filling $E_d - \mu = -\frac{U}{2}$. While in the calculation of the solid line (blue) the full polarization was used, in the calculation of the dashed line (black) we have used the linearized spin susceptibility $\pi\Delta\Pi_0^{\perp}(i\bar{\omega}) \approx 1 - |\bar{\omega}|/\Delta$.

The infinite series of Feynman diagrams which is included in Eq. (2.26) is shown in Fig. 3. Subtracting the Hartree-Fock correction $\delta\xi^{\sigma}$ given in Eq. (2.1) which formally arises from the counterterm in Eq. (2.11), the frequency-dependent part of the self-energy can also be written as

$$\delta\Sigma^{\sigma}(i\omega) \equiv \Sigma^{\sigma}(i\omega) - \delta\xi^{\sigma} = U^2 \int_{\bar{\omega}} \Pi_{\text{LA}}^{\perp}(i\bar{\omega}) G_0^{\bar{\sigma}}(i\omega - i\sigma\bar{\omega}). \quad (2.27)$$

Of particular interest is the quasiparticle residue Z^{σ} , which is defined by

$$Z^{\sigma} = \frac{1}{1 - \left. \frac{\partial \text{Re } \Sigma^{\sigma}(\omega + i0)}{\partial \omega} \right|_{\omega=0}}. \quad (2.28)$$

In Fig. 4 we show the prediction of the LA for the interaction-dependence of the quasiparticle residue in the non-magnetic, particle-hole symmetric case with and without linearization of the spin-flip susceptibility $\Pi_0^{\perp}(i\bar{\omega})$. Obviously, the linearization of the spin-flip susceptibility does lead to modest changes of the Z -factor. The vanishing of Z at $u_0 = 1$ is an unphysical artefact of the LA, which implicitly contains the Stoner instability via the spin susceptibility. In the following section we shall show how to remove this instability by including the

feedback of the fermionic wave function renormalization on the spin susceptibility.

III. FRG WITH PARTIAL BOSONIZATION IN THE SPIN-SINGLET PARTICLE-HOLE CHANNEL

To go beyond the ladder approximation, we now use the collective field FRG approach developed in Refs. [17, 18, 19] to study the mixed Bose-Fermi theory defined by the action $S[\Phi] = S_0[\Phi] + S_1[\Phi]$ in Eqs. (2.10) and (2.11). In fact, this theory has a formal similarity with the theory describing two coupled metallic chains, which was studied by means of the FRG in Ref. [20]. The exact FRG flow equations in the present problem are therefore represented by the same Feynman diagrams as given in Ref. [20].

A. Exact FRG flow equations

To derive exact FRG flow equations, we modify our original model by introducing an infrared cutoff Λ into the Gaussian part (2.10) of the action. In our renormalization group (RG) scheme the cutoff is introduced only in the bosonic sector.^{17,18,19,20} In previous FRG studies of the AIM, fermionic degrees of freedom were directly integrated out and a sharp Matsubara frequency cutoff was employed.^{15,16} For our purpose it is better to work with a smooth cutoff in order to avoid artificial singularities introduced by a sharp cutoff. Formally, we introduce the cutoff via the following substitution in the bosonic part of the Gaussian action $S_0[\Phi]$ given in Eq. (2.10),

$$U^{-1} \rightarrow U^{-1} + R_\Lambda(i\bar{\omega}), \quad (3.1)$$

where

$$R_\Lambda(i\bar{\omega}) = \frac{\Lambda}{\pi\Delta^2} R(|\bar{\omega}|/\Lambda), \quad (3.2)$$

and the function $R(x)$ is given by²¹

$$R(x) = (1-x)\Theta(1-x). \quad (3.3)$$

The flowing spin-flip propagator is then

$$\begin{aligned} F_\Lambda^\perp(i\bar{\omega}) &= [U^{-1} - \Pi_\Lambda^\perp(i\bar{\omega}) + R_\Lambda(i\bar{\omega})]^{-1} \\ &= \frac{U}{1 + U [R_\Lambda(i\bar{\omega}) - \Pi_\Lambda^\perp(i\bar{\omega})]}, \end{aligned} \quad (3.4)$$

where $\Pi_\Lambda^\perp(i\bar{\omega})$ is the flowing irreducible transverse spin susceptibility. Introducing the corresponding single-scale propagator

$$\dot{F}_\Lambda^\perp(i\bar{\omega}) = [-\partial_\Lambda R_\Lambda(i\bar{\omega})][F_\Lambda^\perp(i\bar{\omega})]^2, \quad (3.5)$$




FIG. 5: (Color online) Exact flow equation for the fermionic self-energy $\Sigma_\Lambda^\sigma(\omega)$ with Hubbard-Stratonovich decoupling in the spin-singlet particle-hole channel. In our cutoff scheme only the bosonic propagator is regularized via a cutoff. Thick black arrows denote the flowing fermion propagator given in Eq. (3.8), while thick wavy arrows with a slash denote the flowing single-scale spin-flip propagator defined in Eq. (3.5). The irreducible vertices are denoted by shaded triangles with the appropriate number of external legs. The dots over the fermionic two-point vertices on the left-hand side represent the derivative with respect to the RG cutoff Λ .

the exact FRG flow equation for the irreducible self-energy of spin- σ electrons can then be written as

$$\begin{aligned} \partial_\Lambda \Sigma_\Lambda^\sigma(i\omega) &= \int_{\bar{\omega}} \dot{F}_\Lambda^\perp(i\bar{\omega}) \Gamma_\Lambda^{(\bar{d}_\sigma d_\sigma \bar{\chi} \chi)}(\omega, \omega; \bar{\omega}, \bar{\omega}) \\ &+ \int_{\bar{\omega}} \dot{F}_\Lambda^\perp(i\bar{\omega}) G_\Lambda^\sigma(i\omega - i\sigma\bar{\omega}) \\ &\times \Gamma_\Lambda^{(\bar{d}_\sigma d_\sigma \chi \sigma)}(\omega, \omega - \sigma\bar{\omega}, \sigma\bar{\omega}) \\ &\times \Gamma_\Lambda^{(\bar{d}_\sigma d_\sigma \chi \bar{\sigma})}(\omega - \sigma\bar{\omega}, \omega, \sigma\bar{\omega}). \end{aligned} \quad (3.6)$$

Here, $\Gamma_\Lambda^{(\bar{d}_\sigma d_\sigma \bar{\chi} \chi)}(\omega, \omega; \bar{\omega}, \bar{\omega})$ is the flowing irreducible vertex with two bosonic and two fermionic external legs, and for the labels of the flowing three-legged boson-fermion vertices we have used the short notation $\chi_\uparrow = \chi$ and $\chi_\downarrow = \bar{\chi}$, so that the non-zero combinations are $\Gamma_\Lambda^{(d_\uparrow d_\downarrow \chi_\uparrow)}(\omega, \omega - \bar{\omega}, \bar{\omega}) = \Gamma_\Lambda^{(d_\uparrow d_\downarrow \chi)}(\omega, \omega - \bar{\omega}, \bar{\omega})$ and $\Gamma_\Lambda^{(\bar{d}_\downarrow d_\uparrow \chi_\downarrow)}(\omega - \bar{\omega}, \omega, \bar{\omega}) = \Gamma_\Lambda^{(\bar{d}_\downarrow d_\uparrow \bar{\chi})}(\omega - \bar{\omega}, \omega, \bar{\omega})$. Actually, the three-legged boson-fermion vertices have the symmetry

$$\Gamma_\Lambda^{(\bar{d}_\downarrow d_\uparrow \bar{\chi})}(\omega - \bar{\omega}, \omega, \bar{\omega}) = \Gamma_\Lambda^{(\bar{d}_\uparrow d_\downarrow \chi)}(\omega, \omega - \bar{\omega}, \bar{\omega}), \quad (3.7)$$

so that the two vertices in Eq. (3.6) have the same value. The flowing d -level Green function $G_\Lambda^\sigma(i\omega)$ is related to the flowing self-energy via the Dyson equation,

$$G_\Lambda^\sigma(i\omega) = \frac{1}{i\omega - \xi_0^\sigma - \Delta^\sigma(i\omega) - \Sigma_\Lambda^\sigma(i\omega)}. \quad (3.8)$$

A graphical representation of the exact FRG flow equation (3.6) is shown in Fig. 5. The initial values of the boson-fermion vertices are equal to the bare vertices given in Eq. (2.15), which is unity with our normalization. In principle, the integration of the FRG flow equation (3.6) generates also the Hartree-Fock contribution

(2.3) to the self-energy as a boundary term which is of first-order in U and appears when we integrate this equation up to $\Lambda = 0$. In practice, it is better to drop this first-order term and include the Hartree-Fock self-energy from the beginning into the d -electron propagator, which amounts to imposing the initial condition

$$\Sigma_{\Lambda_0}^\sigma(i\omega) = \delta\xi^\sigma = \frac{U}{2}[n - \sigma m], \quad (3.9)$$

so that the initial $G_{\Lambda_0}^\sigma(i\omega)$ is the Hartree-Fock Green function.

B. Truncation of the FRG equations via Dyson-Schwinger equations

To obtain a closed system of RG flow equations, we need additional RG equations for the four-legged boson-fermion vertex $\Gamma_\Lambda^{(\bar{d}_\sigma d_\sigma \bar{\chi} \chi)}$, for the three-legged boson-fermion vertices $\Gamma_\Lambda^{(\bar{d}_1 d_1 \chi)}$ and $\Gamma_\Lambda^{(\bar{d}_1 d_1 \bar{\chi})}$, as well as for the flowing irreducible spin-flip susceptibility $\Pi_\Lambda^\perp(i\bar{\omega})$ which determines the single-scale spin-flip propagator $\hat{F}_\Lambda^\perp(i\bar{\omega})$. The exact FRG flow equations for the three- and four-legged boson-fermion vertices have been written down diagrammatically in Ref. [20]; the crucial point is that the right-hand sides of these flow equations vanish at the initial scale $\Lambda = \Lambda_0$ because they depend on higher order vertices which are not contained in the bare action. It is therefore reasonable to ignore the RG flow of the three- and four-legged boson-fermion vertices, which amounts to truncating the flow equation (3.6) by replacing these vertices by their initial values,

$$\Gamma_\Lambda^{(\bar{d}_\sigma d_\sigma \bar{\chi} \chi)}(\omega, \omega; \bar{\omega}, \bar{\omega}) \approx \Gamma_{\Lambda_0}^{(\bar{d}_\sigma d_\sigma \bar{\chi} \chi)}(\omega, \omega; \bar{\omega}, \bar{\omega}) = 0, \quad (3.10a)$$

$$\Gamma_\Lambda^{(\bar{d}_1 d_1 \chi)}(\omega, \omega - \bar{\omega}, \bar{\omega}) \approx \Gamma_{\Lambda_0}^{(\bar{d}_1 d_1 \chi)}(\omega, \omega - \bar{\omega}, \bar{\omega}) = 1, \quad (3.10b)$$

$$\Gamma_\Lambda^{(\bar{d}_1 d_1 \bar{\chi})}(\omega - \bar{\omega}, \omega, \bar{\omega}) \approx \Gamma_{\Lambda_0}^{(\bar{d}_1 d_1 \bar{\chi})}(\omega - \bar{\omega}, \omega, \bar{\omega}) = 1, \quad (3.10c)$$

To close our system of flow equations, we still need an additional equation for the flowing spin-flip susceptibility $\Pi_\Lambda^\perp(i\bar{\omega})$, which in our cutoff scheme involves the pure boson vertex with four external legs.²⁰ Fortunately, we can avoid the explicit analysis of this equation by using the Dyson-Schwinger equation for our mixed boson-fermion theory, which implies an exact skeleton equation, relating the flowing spin-flip susceptibility to the flowing fermionic Green function $G_\Lambda^\sigma(i\omega)$ and the flowing three-legged boson-fermion vertices. Using the same method as in Appendix B of Ref. [17], we obtain the skeleton




FIG. 6: (Color online) Graphical representation of the skeleton equation (3.11) relating the flowing spin-flip susceptibility $\Pi_\Lambda^\perp(i\bar{\omega})$ to the exact fermionic Green functions and the three-legged boson-fermion vertices. The two diagrams are equivalent.

equation

$$\begin{aligned} \Pi_\Lambda^\perp(i\bar{\omega}) &= - \int_\omega G_\Lambda^\uparrow(i\omega) G_\Lambda^\downarrow(i\omega - i\bar{\omega}) \\ &\quad \times \Gamma_\Lambda^{(\bar{d}_1 d_1 \chi)}(\omega, \omega - \bar{\omega}, \bar{\omega}) \\ &= - \int_\omega G_\Lambda^\uparrow(i\omega) G_\Lambda^\downarrow(i\omega - i\bar{\omega}) \\ &\quad \times \Gamma_\Lambda^{(\bar{d}_1 d_1 \bar{\chi})}(\omega - \bar{\omega}, \omega, \bar{\omega}). \end{aligned} \quad (3.11)$$

A graphical representation of this equation is shown in Fig. 6. Note that Eq. (3.7) guarantees that the two lines in Eq. (3.11) are indeed identical.

For simplicity, let us focus now on the non-magnetic case, where all correlation functions are spin-independent and we may omit the spin labels. Given our approximation (3.10b), (3.10c), the skeleton equation (3.11) then reduces to

$$\Pi_\Lambda^\perp(i\bar{\omega}) = - \int_\omega G_\Lambda(i\omega) G_\Lambda(i\omega - i\bar{\omega}), \quad (3.12)$$

while the FRG flow equation (3.6) simplifies to

$$\partial_\Lambda \Sigma_\Lambda(i\omega) = \int_{\bar{\omega}} \hat{F}_\Lambda^\perp(i\bar{\omega}) G_\Lambda(i\omega - i\bar{\omega}). \quad (3.13)$$

Eqs. (3.12) and (3.13) form a closed system of integro-differential equations for the flowing self-energy $\Sigma_\Lambda(i\omega)$ of the d -electrons. Recall that Eq. (3.13) depends implicitly on the flowing spin-flip susceptibility via Eqs. (3.4) and (3.5). Anticipating that there is no spontaneous magnetism, the initial condition (3.9) for the fermionic self-energy at scale $\Lambda = \Lambda_0$ is simply

$$\Sigma_{\Lambda_0}(i\omega) = \frac{Un}{2}. \quad (3.14)$$

In the particle-hole symmetric case where $E_d - \mu = -U/2$ and $n = 1$ this cancels precisely the energy $\xi_0^\sigma = E_d - \mu$ in

the Hartree-Fock d -electron propagator $G_{\Lambda_0}(i\omega)$, which in the wide-band limit is therefore given by

$$G_{\Lambda_0}(i\omega) = \frac{1}{i\omega + i\Delta \text{sgn } \omega}. \quad (3.15)$$

We next show that the solution of Eqs. (3.12), (3.13) does not suffer from the Stoner instability and correctly predicts Fermi liquid behavior¹ for arbitrary U . We shall restrict ourselves to the particle-hole symmetric case from now on.

C. Low-energy truncation

The coupled system of integro-differential equations given by Eqs. (3.4), (3.5), (3.12), (3.13) can be solved numerically for arbitrary frequencies, but this is beyond the scope of this work. Here, we shall focus on the low-frequency range $|\omega| \lesssim \Delta$. In this regime, it is reasonable to replace the flowing d -electron propagator appearing on the right-hand sides of Eqs. (3.12), (3.13) by the low-frequency Fermi liquid form

$$G_{\Lambda}(i\omega) = \frac{Z_l}{i\omega + i\Delta_l \text{sgn } \omega}, \quad (3.16)$$

where the flowing wave function renormalization factor is defined by

$$Z_l = \frac{1}{1 - \left. \frac{\partial \Sigma_{\Lambda}(i\omega)}{\partial(i\omega)} \right|_{\omega=0}}, \quad (3.17)$$

and the renormalized flowing hybridization is

$$\Delta_l = Z_l \Delta. \quad (3.18)$$

We consider Z_l and Δ_l to be functions of the logarithmic flow parameter $l = -\ln(\Lambda/\Lambda_0)$. The approximation (3.16) is certainly not sufficient at high frequencies, so that we cannot recover in this way the high-energy Hubbard peaks at strong coupling which are known to appear at an energy scale of order $U/2$.

Given the approximation (3.16), the frequency integration in Eq. (3.12) is easily carried out analytically, resulting in

$$\Pi_{\Lambda}^{\perp}(i\bar{\omega}) = \frac{Z_l^2}{\pi|\bar{\omega}|} \frac{\ln\left(1 + \frac{|\bar{\omega}|}{\Delta_l}\right)}{\left[1 + \frac{|\bar{\omega}|}{2\Delta_l}\right]} = \frac{Z_l}{\pi\Delta} f\left(\frac{|\bar{\omega}|}{\Delta_l}\right), \quad (3.19)$$

with

$$f(x) = \frac{\ln(1+x)}{x(1+x/2)}. \quad (3.20)$$

Introducing the dimensionless coupling constants

$$u_l = Z_l u_0 = \frac{Z_l U}{\pi\Delta} = \frac{Z_l^2 U}{\pi\Delta_l}, \quad (3.21)$$

$$\lambda_l = \frac{\Lambda}{Z_l \Delta} = \frac{\Lambda}{\Delta_l}, \quad (3.22)$$

the single-scale propagator defined in (3.5) can then be written as

$$\dot{F}_{\Lambda}^{\perp}(i\bar{\omega}) = -\frac{\pi}{Z_l^2} \frac{\Theta(\Lambda - |\bar{\omega}|)}{\left[\frac{1}{u_l} + \lambda_l - \frac{|\bar{\omega}|}{\Delta_l} - f\left(\frac{|\bar{\omega}|}{\Delta_l}\right)\right]^2}. \quad (3.23)$$

Actually, to be consistent with our approximation (3.16) which is based on the expansion of the fermionic self-energy to linear order in frequency, we should also expand the bosonic self-energy $\Pi_{\Lambda}^{\perp}(i\bar{\omega})$ to linear order in $\bar{\omega}$. Using $f(x) = 1 - x + \mathcal{O}(x^2)$, we have

$$\pi\Delta\Pi_{\Lambda}^{\perp}(i\bar{\omega}) \approx Z_l - \frac{|\bar{\omega}|}{\Delta} + \mathcal{O}(\bar{\omega}^2). \quad (3.24)$$

As compared with the non-interacting susceptibility (2.20), the leading constant term in Eq. (3.24) is reduced by the wave function renormalization factor, while the linear term is not renormalized. In this approximation, our single-scale propagator (3.23) simplifies to

$$\dot{F}_{\Lambda}^{\perp}(i\bar{\omega}) = -\frac{\pi}{Z_l^2} \frac{\Theta(\Lambda - |\bar{\omega}|)}{\left[\frac{1}{u_l} + \lambda_l - 1\right]^2}, \quad (3.25)$$

which by construction of the regulator function given in Eqs. (3.2) and (3.3) is constant for frequencies below the running cutoff Λ . To determine Z_l , we calculate the flowing anomalous dimension η_l , which is directly related to the derivative of the self-energy with respect to the flow parameter,

$$\eta_l = -\partial_l \ln Z_l = Z_l \Lambda \lim_{\omega \rightarrow 0} \frac{\partial}{\partial(i\omega)} \partial_{\Lambda} \Sigma_{\Lambda}(i\omega). \quad (3.26)$$

Substituting Eqs. (3.13) and (3.16), we obtain

$$\eta_l = Z_l^2 \Lambda \lim_{\omega \rightarrow 0} \frac{\partial}{\partial(i\omega)} \int_{\bar{\omega}} \frac{\dot{F}_{\Lambda}^{\perp}(i\bar{\omega})}{i\omega - i\bar{\omega} + i\Delta_l \text{sgn}(\omega - \bar{\omega})}. \quad (3.27)$$

If we naïvely interchange the order of integration and differentiation, we encounter an ambiguous expression of the form

$$\begin{aligned} & \lim_{\omega \rightarrow 0} \frac{\partial}{\partial(i\omega)} \left[\frac{1}{i\omega - i\bar{\omega} + i\Delta_l \text{sgn}(\omega - \bar{\omega})} \right] \\ &= \frac{1}{[\bar{\omega} + \Delta_l \text{sgn } \bar{\omega}]^2} + \frac{2\Delta_l \delta(\bar{\omega})}{[\Delta_l (2\Theta(\bar{\omega}) - 1)]^2}, \end{aligned} \quad (3.28)$$

where we have written $\text{sgn } \bar{\omega} = 2\Theta(\bar{\omega}) - 1$. As pointed out by Morris,²² one should interpret the product of the delta function $\delta(x)$ with any function $f(\Theta(x))$ of the step function as

$$\delta(x)f(\Theta(x)) = \delta(x) \int_0^1 dt f(t). \quad (3.29)$$

Using this relation, Eq. (3.28) reduces to

$$\begin{aligned} & \lim_{\omega \rightarrow 0} \frac{\partial}{\partial(i\omega)} \left[\frac{1}{i\omega - i\bar{\omega} + i\Delta_l \text{sgn}(\omega - \bar{\omega})} \right] \\ &= \frac{1}{[|\bar{\omega}| + \Delta_l]^2} - \frac{2\delta(\bar{\omega})}{\Delta_l}, \end{aligned} \quad (3.30)$$




FIG. 7: Numerical solution of Eq. (3.33) for different values of the bare coupling $u_0 = U/(\pi\Delta)$. Here, l_* is the scale where the running cutoff Λ is equal to the hybridization Δ , i.e., $\Lambda_0 e^{-l_*} = \Delta$.

so that Eq. (3.27) takes the form

$$\eta_l = -\frac{Z_l \Lambda}{\pi \Delta} \dot{F}_\Lambda^\perp(i0) + Z_l^2 \Lambda \int_{\bar{\omega}} \frac{\dot{F}_\Lambda^\perp(i\bar{\omega})}{(|\bar{\omega}| + \Delta_l)^2}. \quad (3.31)$$

With the single-scale propagator given in Eq. (3.25), the frequency integration is trivial and we finally obtain from Eq. (3.27) for the flowing anomalous dimension

$$\eta_l = \frac{\lambda_l}{[1 + \lambda_l][\frac{1}{u_l} + \lambda_l - 1]^2}. \quad (3.32)$$

The right-hand side of this expression depends on Z_l via the flowing couplings u_l and λ_l defined in Eqs. (3.21) and (3.22) so that

$$\partial_l Z_l = -\eta_l Z_l \quad (3.33)$$

is an ordinary differential equation for the flowing wave function renormalization factor Z_l , which is easily solved numerically. The result for the flowing Z_l is shown in Fig. 7. Obviously, for $l \rightarrow \infty$ the wave function renormalization factor approaches a finite limit, $Z = \lim_{l \rightarrow \infty} Z_l$, which we show in Fig. 8 as a function of the bare coupling u_0 . In contrast to the ladder approximation discussed in Sec. II we now obtain a finite Z for all values of the interaction, so that the fluctuations included in our simple FRG approach are sufficient to remove the unphysical Stoner instability. On the other hand, quantitatively our truncation of the exact FRG flow equations does not reproduce the correct strong-coupling behavior of the quasiparticle residue, which is known to exhibit the same exponential suppression as the Kondo temperature for $u_0 \rightarrow \infty$ (see Ref. [23]),

$$Z_{\text{exact}} \sim \sqrt{\frac{8u_0}{\pi}} \exp[-\pi^2 u_0/8]. \quad (3.34)$$

From the numerical solution of our flow equation given in Eqs. (3.32), (3.33) we find asymptotically

$$Z \sim \frac{0.445}{u_0}, \quad u_0 \gg 1. \quad (3.35)$$




FIG. 8: The solid line is the wave function renormalization factor $Z = \lim_{l \rightarrow \infty} Z_l$ as a function of the bare coupling u_0 obtained from the numerical solution of our truncated flow equation (3.33). For comparison, we show as the dashed line the result of the ladder approximation.

The fact that this is much larger than the exact result (3.34) indicates that our approach underestimates the strength of fluctuations. In the following section we shall therefore include the longitudinal spin fluctuation channel and show that it indeed further suppresses the value of Z , although within our approximations we are unable to recover the exponential suppression described by Eq. (3.34).

IV. FRG WITH PARTIAL BOSONIZATION OF TRANSVERSE AND LONGITUDINAL SPIN FLUCTUATIONS

A. Multi-channel Hubbard-Stratonovich transformation

Using the antisymmetry of the Grassmann-fields, the local Hubbard interaction can be written in infinitely many equivalent ways, such as

$$U n_\uparrow(\tau) n_\downarrow(\tau) = -\frac{U^\parallel}{2} m^2(\tau) - U^\perp \bar{s}(\tau) s(\tau), \quad (4.1)$$

where

$$U^\parallel + U^\perp = U. \quad (4.2)$$

Here, $n_\sigma(\tau) = \bar{d}_\sigma(\tau) d_\sigma(\tau)$ and the composite fields $m(\tau) = \sum_\sigma \sigma n_\sigma(\tau)$, $\bar{s}(\tau) = \bar{d}_\uparrow(\tau) d_\downarrow(\tau)$, and $s(\tau) = \bar{d}_\downarrow(\tau) d_\uparrow(\tau)$ represent the longitudinal and transverse spin components, see also Eq. (2.8). Note that by construction of the decomposition (4.1) we have included both longitudinal and transverse spin fluctuations, but no charge fluctuations. In Sec. III we have satisfied Eq. (4.1) by setting $U^\parallel = 0$ and $U^\perp = U$, which is the natural choice if one is interested in bosonizing the transverse spin fluctuations. Alternatively, one could set $U^\parallel = U$ and $U^\perp = 0$, so that longitudinal spin fluctuations can be introduced

via a suitable Hubbard-Stratonovich field. The ambiguity associated with Eqs. (4.1) and (4.2) has been discussed for many decades in the literature.^{24,25,26,27,28,29} Depending on the physical problem of interest, certain special choices of $U^\parallel$ and $U^\perp$ can be advantageous.²⁸ At this point, we simply leave the precise values of $U^\parallel$ and $U^\perp$ unspecified but assume that both are positive and satisfy Eq. (4.2). Later we shall show that in the strong coupling regime of the particle-hole symmetric AIM the optimal choice is $U^\parallel = U/3$ and $U^\perp = 2U/3$. In this case the decomposition (4.1) is manifestly spin-rotational invariant and can be written as

$$Un_\uparrow(\tau)n_\downarrow(\tau) = -\frac{U}{6}(\vec{s}(\tau))^2, \quad (4.3)$$

where the composite vector field $\vec{s}(\tau) = d^\dagger(\tau)\vec{\sigma}d(\tau)$ represents the spin vector. Here, $d^\dagger(\tau) = [\bar{d}_\uparrow(\tau), \bar{d}_\downarrow(\tau)]$ and $\vec{\sigma}$ is the usual matrix vector of Pauli matrices.

Starting from the representation (4.1), we decouple the interaction in the longitudinal spin channel using a real Hubbard-Stratonovich field η and in the transverse spin channel using the complex Hubbard-Stratonovich fields χ and $\bar{\chi}$ introduced in Sec. II B. The partition function can then be written as in Eq. (2.9), where $\Phi = [d_\uparrow, \bar{d}_\uparrow, d_\downarrow, \bar{d}_\downarrow, \eta, \chi, \bar{\chi}]$ is now a seven-component superfield. For the Gaussian part of the action we then obtain instead of Eq. (2.10)

$$S_0[\Phi] = - \int_\omega \sum_\sigma [G_0^\sigma(i\omega)]^{-1} \bar{d}_{\omega\sigma} d_{\omega\sigma} + \int_{\bar{\omega}} (U^\perp)^{-1} \bar{\chi}_{\bar{\omega}} \chi_{\bar{\omega}} + \frac{1}{2} \int_{\bar{\omega}} (U^\parallel)^{-1} \eta_{-\bar{\omega}} \eta_{\bar{\omega}}, \quad (4.4)$$

and the interaction part can be written as

$$S_1[\Phi] = \int_{\bar{\omega}} [\bar{s}_{\bar{\omega}} \chi_{\bar{\omega}} + s_{\bar{\omega}} \bar{\chi}_{\bar{\omega}} + m_{-\bar{\omega}} \eta_{\bar{\omega}}] - \int_\omega \sum_\sigma \delta\xi^\sigma \bar{d}_{\omega\sigma} d_{\omega\sigma}. \quad (4.5)$$

The counterterm $\delta\xi^\sigma$ in Eq. (4.5) is subtracted to correct for the inclusion of the Hartree-Fock self-energy in the propagator $G_0^\sigma(i\omega)$ in Eq. (4.4). While the transverse spin components $s_{\bar{\omega}}$ and $\bar{s}_{\bar{\omega}}$ are given by Eqs. (2.12) and (2.13), the longitudinal spin component is given by

$$m_{\bar{\omega}} = \int_0^\beta d\tau e^{i\bar{\omega}\tau} m(\tau) = \int_\omega \sum_\sigma \sigma \bar{d}_{\omega\sigma} d_{\omega+\bar{\omega},\sigma}. \quad (4.6)$$

For simplicity, we shall focus here on the non-magnetic, particle-hole-symmetric case. Since in the strong coupling limit the low-energy physics is expected to be dominated by spin fluctuations, we set up the FRG by introducing a cutoff only in the bosonic fields η, χ and $\bar{\chi}$ associated with spin fluctuations. Particle-hole symmetry guarantees that the exact self-energy at zero frequency cancels the term $\epsilon_d - \mu$, so that we may approximate $G_0^\sigma(i\omega)$ by Eq. (3.15).

B. FRG flow equations

We now generalize Eq. (3.1) by introducing cutoffs into both types of bosonic propagators appearing in the Gaussian action (4.4),

$$(U^\alpha)^{-1} \rightarrow (U^\alpha)^{-1} + R_\Lambda^\alpha(i\bar{\omega}), \quad (4.7)$$

where $\alpha = \parallel, \perp$ labels the two types of spin fluctuations, and the cutoff functions are

$$R^\parallel(i\bar{\omega}) = \frac{2\Lambda}{\pi\Delta^2} R\left(\frac{|\bar{\omega}|}{\Lambda}\right), \quad (4.8a)$$

$$R^\perp(i\bar{\omega}) = \frac{\Lambda}{\pi\Delta^2} R\left(\frac{|\bar{\omega}|}{\Lambda}\right), \quad (4.8b)$$

with $R(x)$ given in Eq. (3.3). From the general FRG flow equations for mixed boson-fermion theories given in Refs. [17,19] we then obtain the following exact flow equation for the self-energy,

$$\begin{aligned} \partial_\Lambda \Sigma_\Lambda^\sigma(i\omega) &= \int_{\bar{\omega}} \dot{F}_\Lambda^\parallel(i\bar{\omega}) \Gamma_\Lambda^{(\bar{d}_\sigma d_\sigma \eta \eta)}(\omega, \omega; \bar{\omega}, -\bar{\omega}) + \int_{\bar{\omega}} \dot{F}_\Lambda^\perp(i\bar{\omega}) \Gamma_\Lambda^{(\bar{d}_\sigma d_\sigma \bar{\chi} \chi)}(\omega, \omega; \bar{\omega}, \bar{\omega}) \\ &+ \int_{\bar{\omega}} \dot{F}_\Lambda^\parallel(i\bar{\omega}) G_\Lambda^\sigma(i\omega - i\bar{\omega}) \Gamma_\Lambda^{(\bar{d}_\sigma d_\sigma \eta)}(\omega, \omega - \bar{\omega}, \bar{\omega}) \Gamma_\Lambda^{(\bar{d}_\sigma d_\sigma \eta)}(\omega - \bar{\omega}, \omega, -\bar{\omega}) \\ &+ \int_{\bar{\omega}} \dot{F}_\Lambda^\perp(i\bar{\omega}) G_\Lambda^\sigma(i\omega - i\sigma\bar{\omega}) \Gamma_\Lambda^{(\bar{d}_\sigma d_\sigma \chi \sigma)}(\omega, \omega - \sigma\bar{\omega}, \sigma\bar{\omega}) \Gamma_\Lambda^{(\bar{d}_\sigma d_\sigma \chi \sigma)}(\omega - \sigma\bar{\omega}, \omega, \sigma\bar{\omega}), \end{aligned} \quad (4.9)$$

which generalizes Eq. (3.6) and is shown graphically in Fig. 9. The superscripts label the different types of vertices; for the boson-fermion vertices associated with the transverse spin field we have used again the short nota-

tion $\chi_\uparrow = \chi$ and $\chi_\downarrow = \bar{\chi}$, see Eq. (3.6). The longitudinal and transverse single-scale propagators are

$$\dot{F}_\Lambda^\alpha(i\bar{\omega}) = [-\partial_\Lambda R_\Lambda^\alpha(i\bar{\omega})][F_\Lambda^\alpha(i\bar{\omega})]^2, \quad (4.10)$$




FIG. 9: (Color online) Graphical representation of the exact FRG flow equation (4.9) for the self-energies $\Sigma_\Lambda^\uparrow$ and $\Sigma_\Lambda^\downarrow$. The longitudinal spin-field η is represented by dashed lines; the other symbols are the same as in Fig. 5.

with

$$F_\Lambda^\alpha(i\bar{\omega}) = [(U^\alpha)^{-1} - \Pi_\Lambda^\alpha(i\bar{\omega}) + R_\Lambda^\alpha(i\bar{\omega})]^{-1}, \quad (4.11)$$

where $\Pi_\Lambda^\alpha(i\bar{\omega})$ are the corresponding irreducible spin susceptibilities. As discussed in Sec. III B, we use the skeleton equations (which can be formally derived from the Dyson-Schwinger equations) to relate the flow of the susceptibilities to the flow of the d -electron propagators and the three-legged boson-fermion vertex. The skeleton equation for the transverse susceptibility $\Pi_\Lambda^\perp(i\bar{\omega})$ has already been discussed in Sec. III B, see Eq. (3.11); it can also be written as

$$\begin{aligned} \Pi_\Lambda^\perp(i\bar{\omega}) = & - \int_{\omega} G_\Lambda^\sigma(i\omega) G_\Lambda^{\bar{\sigma}}(i\omega - i\sigma\bar{\omega}) \\ & \times \Gamma_\Lambda^{(\bar{d}_\sigma d_{\bar{\sigma}} \chi_\sigma)}(\omega, \omega - \sigma\bar{\omega}, \bar{\omega}). \end{aligned} \quad (4.12)$$

The corresponding skeleton equation for the longitudinal spin susceptibility is

$$\begin{aligned} \Pi_\Lambda^\parallel(i\bar{\omega}) = & - \int_{\omega} \sum_{\sigma} \sigma G_\Lambda^\sigma(i\omega) G_\Lambda^\sigma(i\omega - i\bar{\omega}) \\ & \times \Gamma_\Lambda^{(\bar{d}_\sigma d_\sigma \eta)}(\omega, \omega - \bar{\omega}, \bar{\omega}). \end{aligned} \quad (4.13)$$

To obtain a closed RG flow equation for the self-energy, we still need flow equations for the vertices with three and four external legs appearing in Eq. (4.9). As in Sec. III B, we simply set all mixed boson-fermion vertices with four external legs equal to zero, because these vertices vanish at the initial scale. However, in contrast to the FRG with only transverse spin fluctuations discussed in Sec. III B, the renormalization of the three-legged boson-fermion vertices is now important. In the

approximation where mixed boson-fermion vertices with four and more external legs are ignored, the FRG flow equation for the longitudinal three-legged boson-fermion vertex is

$$\begin{aligned} & \partial_\Lambda \Gamma_\Lambda^{(\bar{d}_\sigma d_\sigma \eta)}(\omega + \bar{\omega}, \omega, \bar{\omega}) \\ = & \int_{\bar{\omega}'} \dot{F}_\Lambda^\parallel(i\bar{\omega}') G_\Lambda^\sigma(i\omega + i\bar{\omega} + i\bar{\omega}') G_\Lambda^\sigma(i\omega + i\bar{\omega}') \\ & \times \Gamma_\Lambda^{(\bar{d}_\sigma d_\sigma \eta)}(\omega + \bar{\omega}, \omega + \bar{\omega} + \bar{\omega}', -\bar{\omega}') \\ & \times \Gamma_\Lambda^{(\bar{d}_\sigma d_\sigma \eta)}(\omega + \bar{\omega} + \bar{\omega}', \omega + \bar{\omega}', \bar{\omega}) \\ & \times \Gamma_\Lambda^{(\bar{d}_\sigma d_\sigma \eta)}(\omega + \bar{\omega}', \omega, \bar{\omega}') \\ + & \int_{\bar{\omega}'} \dot{F}_\Lambda^\perp(-i\sigma\bar{\omega}') G_\Lambda^{\bar{\sigma}}(i\omega + i\bar{\omega} + i\bar{\omega}') G_\Lambda^{\bar{\sigma}}(i\omega + i\bar{\omega}') \\ & \times \Gamma_\Lambda^{(\bar{d}_\sigma d_{\bar{\sigma}} \chi_\sigma)}(\omega + \bar{\omega}, \omega + \bar{\omega} + \bar{\omega}', -\sigma\bar{\omega}') \\ & \times \Gamma_\Lambda^{(\bar{d}_\sigma d_\sigma \eta)}(\omega + \bar{\omega} + \bar{\omega}', \omega + \bar{\omega}', \bar{\omega}) \\ & \times \Gamma_\Lambda^{(\bar{d}_{\bar{\sigma}} d_\sigma \chi_{\bar{\sigma}})}(\omega + \bar{\omega}', \omega, -\sigma\bar{\omega}'), \end{aligned} \quad (4.14)$$

while the corresponding flow equation for the transverse vertex reads

$$\begin{aligned} & \partial_\Lambda \Gamma_\Lambda^{(\bar{d}_\sigma d_{\bar{\sigma}} \chi_\sigma)}(\omega + \bar{\omega}, \omega, \sigma\bar{\omega}) \\ = & \int_{\bar{\omega}'} \dot{F}_\Lambda^\parallel(i\bar{\omega}') G_\Lambda^\sigma(i\omega + i\bar{\omega} + i\bar{\omega}') G_\Lambda^{\bar{\sigma}}(i\omega + i\bar{\omega}') \\ & \times \Gamma_\Lambda^{(\bar{d}_\sigma d_\sigma \eta)}(\omega + \bar{\omega}, \omega + \bar{\omega} + \bar{\omega}', -\bar{\omega}') \\ & \times \Gamma_\Lambda^{(\bar{d}_\sigma d_{\bar{\sigma}} \chi_\sigma)}(\omega + \bar{\omega} + \bar{\omega}', \omega + \bar{\omega}', \sigma\bar{\omega}) \\ & \times \Gamma_\Lambda^{(\bar{d}_{\bar{\sigma}} d_\sigma \eta)}(\omega + \bar{\omega}', \omega, \bar{\omega}'). \end{aligned} \quad (4.15)$$

Graphical representations of Eqs. (4.14) and (4.15) are shown in Figs. 10 and 11. From $S_1[\Phi]$ in Eq. (4.5) we see that the initial conditions for the three-legged vertices at scale $\Lambda = \Lambda_0$ are

$$\Gamma_{\Lambda_0}^{(\bar{d}_\sigma d_\sigma \eta)}(\omega + \bar{\omega}, \omega, \bar{\omega}) = \sigma, \quad (4.16a)$$

$$\Gamma_{\Lambda_0}^{(\bar{d}_\sigma d_{\bar{\sigma}} \chi_\sigma)}(\omega + \bar{\omega}, \omega, \sigma\bar{\omega}) = 1. \quad (4.16b)$$

C. Low-energy truncation

To make further progress, we now neglect the frequency-dependence of the three-legged boson-fermion vertices, setting

$$\Gamma_\Lambda^{(\bar{d}_\sigma d_\sigma \eta)}(\omega + \bar{\omega}, \omega, \bar{\omega}) \approx \sigma \gamma_\Lambda^\parallel, \quad (4.17)$$

$$\Gamma_\Lambda^{(\bar{d}_\sigma d_{\bar{\sigma}} \chi_\sigma)}(\omega + \bar{\omega}, \omega, \sigma\bar{\omega}) \approx \gamma_\Lambda^\perp. \quad (4.18)$$

Keeping in mind that we have also neglected all vertices involving more than three external legs, our exact FRG flow equation (4.9) for the self-energy reduces to

$$\begin{aligned} \partial_\Lambda \Sigma_\Lambda^\sigma(i\omega) = & (\gamma_\Lambda^\parallel)^2 \int_{\bar{\omega}} \dot{F}_\Lambda^\parallel(i\bar{\omega}) G_\Lambda^\sigma(i\omega - i\bar{\omega}) \\ & + (\gamma_\Lambda^\perp)^2 \int_{\bar{\omega}} \dot{F}_\Lambda^\perp(i\bar{\omega}) G_\Lambda^{\bar{\sigma}}(i\omega - i\sigma\bar{\omega}). \end{aligned} \quad (4.19)$$




FIG. 10: (Color online) Graphical representation of the FRG flow equation (4.14) for the longitudinal boson-fermion vertices $\Gamma_\Lambda^{(\bar{d}_\uparrow d_\uparrow \eta)}$ (upper graph) and $\Gamma_\Lambda^{(\bar{d}_\downarrow d_\downarrow \eta)}$ (lower graph).




FIG. 11: (Color online) Graphical representation of the FRG flow equation (4.15) for the transverse boson-fermion vertices $\Gamma_\Lambda^{(\bar{d}_\uparrow d_\perp \chi)}$ (upper graph) and $\Gamma_\Lambda^{(\bar{d}_\downarrow d_\perp \chi)}$ (lower graph).

Moreover, the approximate FRG flow equation (4.14) for the longitudinal spin-fermion vertex reduces to

$$\begin{aligned} \partial_\Lambda \gamma_\Lambda^\parallel &= (\gamma_\Lambda^\parallel)^3 \int_{\bar{\omega}} \dot{F}_\Lambda^\parallel(i\bar{\omega}) [G_\Lambda^\dagger(i\bar{\omega})]^2 \\ &- \gamma_\Lambda^\parallel (\gamma_\Lambda^\perp)^2 \int_{\bar{\omega}} \dot{F}_\Lambda^\perp(i\bar{\omega}) [G_\Lambda^\dagger(-i\bar{\omega})]^2, \end{aligned} \quad (4.20)$$

while the FRG equation (4.15) for the transverse spin-fermion vertex becomes

$$\partial_\Lambda \gamma_\Lambda^\perp = -\gamma_\Lambda^\perp (\gamma_\Lambda^\parallel)^2 \int_{\bar{\omega}} \dot{F}_\Lambda^\parallel(i\bar{\omega}) G_\Lambda^\dagger(i\bar{\omega}) G_\Lambda^\perp(i\bar{\omega}). \quad (4.21)$$

Within the same approximation, the skeleton equations (4.12) and (4.13) for the spin susceptibilities are

$$\Pi_\Lambda^\parallel(i\bar{\omega}) = -\gamma_\Lambda^\parallel \int_{\omega} \sum_{\sigma} G_\Lambda^\sigma(i\omega) G_\Lambda^\sigma(i\omega - i\bar{\omega}), \quad (4.22)$$

$$\Pi_\Lambda^\perp(i\bar{\omega}) = -\gamma_\Lambda^\perp \int_{\omega} G_\Lambda^\sigma(i\omega) G_\Lambda^{\bar{\sigma}}(i\omega - i\sigma\bar{\omega}). \quad (4.23)$$

Eqs. (4.19)–(4.23) form a closed system of integro-differential equations for the frequency-dependent self-energy $\Sigma_\Lambda^\sigma(i\omega)$ and the frequency-independent parts $\gamma_\Lambda^\parallel$ and $\gamma_\Lambda^\perp$ of the three-legged boson-fermion vertices. In order to make progress analytically, let us now approximate the flowing d -electron Green function on the right-hand sides of the flow equation by its low-energy Fermi liquid form (3.16). Then the integrations in Eqs. (4.22), (4.23) can be performed analytically. For simplicity, we focus again on the non-magnetic particle-hole symmetric case and omit the spin-labels. We then obtain

$$\Pi_\Lambda^\parallel(i\bar{\omega}) = \frac{2Z_l}{\pi\Delta} \gamma_l^\parallel f\left(\frac{|\bar{\omega}|}{\Delta_l}\right), \quad (4.24)$$

$$\Pi_\Lambda^\perp(i\bar{\omega}) = \frac{Z_l}{\pi\Delta} \gamma_l^\perp f\left(\frac{|\bar{\omega}|}{\Delta_l}\right), \quad (4.25)$$

where the function $f(x)$ is given in Eq. (3.20). The bosonic single-scale propagators can then be written as

$$\dot{F}_\Lambda^\parallel(i\bar{\omega}) = -\frac{\pi}{2Z_l^2} \frac{\Theta(\Lambda - |\bar{\omega}|)}{\left[\frac{1}{u_l^\parallel} + \lambda_l - \frac{|\bar{\omega}|}{\Delta_l} - \gamma_l^\parallel f\left(\frac{|\bar{\omega}|}{\Delta_l}\right)\right]^2}, \quad (4.26)$$

$$\dot{F}_\Lambda^\perp(i\bar{\omega}) = -\frac{\pi}{Z_l^2} \frac{\Theta(\Lambda - |\bar{\omega}|)}{\left[\frac{1}{u_l^\perp} + \lambda_l - \frac{|\bar{\omega}|}{\Delta_l} - \gamma_l^\perp f\left(\frac{|\bar{\omega}|}{\Delta_l}\right)\right]^2}. \quad (4.27)$$

Here, we have introduced again the notation $\lambda_l = \Lambda/\Delta_l = \Lambda/(Z_l\Delta)$, and the running interaction constants $u_l^\parallel$ and $u_l^\perp$ are defined by

$$u_l^\parallel = Z_l \frac{2U^\parallel}{\pi\Delta}, \quad u_l^\perp = Z_l \frac{U^\perp}{\pi\Delta}. \quad (4.28)$$

In terms of the dimensionless bare couplings $u_0^\parallel$ and $u_0^\perp$ the condition (4.2) reads

$$u_0^\parallel/2 + u_0^\perp = u_0. \quad (4.29)$$

As discussed in Sec. III C, for consistency we expand the function $f(|\bar{\omega}|/\Delta_l)$ in Eqs. (4.24) and (4.25) to linear order in frequency as in Eq. (3.24). Then the frequency integrations in Eqs. (4.19), (4.20), (4.21) can be performed analytically and we obtain the following system of RG flow equations for the three running couplings Z_l , $\gamma_l^\parallel$,

and $\gamma_l^\perp$,

$$\frac{\partial_l Z_l}{Z_l} = -\eta_l^\parallel - \eta_l^\perp, \quad (4.30)$$

$$\frac{\partial_l \gamma_l^\parallel}{\gamma_l^\parallel} = -\frac{1}{2}A_l^\parallel + A_l^\perp, \quad (4.31)$$

$$\frac{\partial_l \gamma_l^\perp}{\gamma_l^\perp} = \frac{1}{2}A_l^\parallel, \quad (4.32)$$

where

$$\eta_l^\parallel = \frac{(\gamma_l^\parallel)^2 \lambda_l}{2[1/u_l^\parallel + \lambda_l - \gamma_l^\parallel]^2} - \frac{1}{2}A_l^\parallel, \quad (4.33)$$

$$\eta_l^\perp = \frac{(\gamma_l^\perp)^2 \lambda_l}{[1/u_l^\perp + \lambda_l - \gamma_l^\perp]^2} - A_l^\perp, \quad (4.34)$$

$$A_l^\alpha = (\gamma_l^\alpha)^2 \lambda_l I(1/u_l^\alpha + \lambda_l - \gamma_l^\alpha, \gamma_l^\alpha - 1, \lambda_l). \quad (4.35)$$

Here, we have introduced the dimensionless integral

$$\begin{aligned} I(a, b, \lambda) &= \int_0^\lambda dx \frac{1}{(a + bx)^2(1+x)^2} \\ &= \frac{\lambda}{(a-b)^2} \left[\frac{1}{1+\lambda} + \frac{b^2}{a(a+b\lambda)} \right] \\ &\quad - \frac{2b}{(a-b)^3} \ln \left[\frac{a(1+\lambda)}{a+b\lambda} \right]. \end{aligned} \quad (4.36)$$

Of special interest is the manifestly spin-rotationally invariant decomposition (4.3). In this case $U^\parallel = U^\perp/2 = U/3$, such that each of the longitudinal and two transverse channels has the same weight. In terms of the dimensionless bare couplings $u_0^\parallel$ and $u_0^\perp$ defined in Eq. (4.28) we therefore have $u_0^\parallel = u_0^\perp = 2u_0/3$. It is now easy to see that for any l we have $u_l^\parallel = u_l^\perp = 2u_l/3$ with $u_l = Z_l u_0$ and $\gamma_l \equiv \gamma_l^\parallel = \gamma_l^\perp$. Our flow equations (4.30)–(4.32) then reduce to

$$\frac{\partial_l Z_l}{Z_l} = -\eta_l, \quad (4.37)$$

$$\frac{\partial_l \gamma_l}{\gamma_l} = \frac{1}{2}A_l, \quad (4.38)$$

where

$$\eta_l = \frac{3\gamma_l^2 \lambda_l}{2[3/(2u_l) + \lambda_l - \gamma_l]^2} - \frac{3}{2}A_l, \quad (4.39)$$

$$A_l = \gamma_l^2 \lambda_l I(3/(2u_l) + \lambda_l - \gamma_l, \gamma_l - 1, \lambda_l). \quad (4.40)$$

The factors of 3 appearing here are related to the fact that for this choice of parameters we have three equivalent channels. Clearly, our truncated flow equations respect the spin-rotation symmetry of the problem such that for this special decomposition we can expect the best results.




FIG. 12: (Color online) Typical flow of the wave function renormalization factor Z_l and of the frequency-independent parts $\gamma_l^\parallel$ and $\gamma_l^\perp$ of the boson-fermion vertices obtained from the numerical solution of Eqs. (4.30)–(4.32). The scale $l_* = \ln(\Lambda_0/\Delta)$ is the same as in Fig. 7. The curves are for $u_0^\parallel = 1$ and $u_0^\perp = 1.5$, such that $u_0 = u_0^\parallel/2 + u_0^\perp = 2$.

D. Results

The three-dimensional system of differential equations given in Eqs. (4.30)–(4.32) can easily be solved numerically. An example of a typical RG flow is shown in Fig. 12. Obviously, the wave function renormalization factor decreases monotonically as the RG is iterated, while the vertex corrections $\gamma_l^\parallel$ and $\gamma_l^\perp$ both increase. The strongest variations occur at the scale l_* where the effective cutoff $\Lambda_0 e^{-l_*}$ is equal to the hybridization Δ .

Before discussing the behavior of $Z = \lim_{l \rightarrow \infty} Z_l$ in the strong coupling regime, let us fix the optimal choice of the relative weight of the bare couplings $U^\parallel$ and $U^\perp$ in the Hubbard-Stratonovich decoupling (4.1). In Fig. 13 we show the dependence of the wave function renormalization factor on the choice of $u_0^\parallel/u_0^\perp$ for fixed values of u_0 . It turns out that all curves have a pronounced plateau with a minimum precisely at $u_0^\parallel/u_0^\perp = 1$, which corresponds to the manifestly spin-rotationally invariant decoupling (4.3). If we fix the choice of the Hubbard-Stratonovich decoupling by demanding that first-order variations of the results around the optimal choice should vanish, then we are naturally led to the decoupling (4.3). For the rest of this section we shall therefore use the manifestly spin-rotationally invariant choice $u_0^\parallel = u_0^\perp = 2u_0/3$.

In Fig. 14 we show our numerical results for Z as a function of the dimensionless bare coupling u_0 . For comparison, we also show the prediction of the ladder approximation, as well as our FRG results with decoupling only in the transverse spin channel, and numerically accurate results obtained within Wilson's numerical renormalization group.¹⁶ Our FRG calculation with simultaneous decoupling in both transverse and longitudinal spin channels obviously yields much better results for the suppression of Z in the strong coupling regime than the single-




FIG. 13: (Color online) Dependence of the quasiparticle residue Z on the choice of the Hubbard-Stratonovich decoupling, parameterized by the ratio $u_0^{\parallel}/u_0^{\perp}$, with $u_0^{\parallel}/2 + u_0^{\perp} = u_0$. All curves exhibit a local minimum precisely at $u_0^{\parallel}/u_0^{\perp} = 1$, corresponding to the manifestly spin-rotationally invariant decoupling $u_0^{\parallel} = u_0^{\perp} = 2u_0/3$. While this minimum is also the global minimum for $u_0 = 1, 2$ and 5 , it is only a local minimum for $u_0 = 0.5$. In fact, for $u_0 = 0.5$ the curve is almost flat and has a global minimum at $u_0^{\parallel}/u_0^{\perp} = 0$ and a local maximum between the two minima.




FIG. 14: (Color online) Wave function renormalization factor $Z = \lim_{l \rightarrow \infty} Z_l$ as a function of the bare coupling $u_0 = U/(\pi\Delta)$ obtained from the numerical solution of Eqs. (4.30)–(4.32). The two-channel FRG results are for the manifestly spin-rotationally invariant choice of the two Hubbard-Stratonovich decouplings where $u_0^{\parallel} = u_0^{\perp} = 2u_0/3$. For a comparison, we have also shown NRG results from Ref. [16] (which we have extrapolated to values $u_0 \gtrsim 4.5$).

channel FRG discussed in Sec. III. In fact, on the scale of Fig. 14 our FRG results for Z seem to be reasonably close to the exact numerical results, which in the strong coupling regime can be approximated by the asymptotic formula (3.34). To investigate whether our FRG approach reproduces the known exponential suppression of Z , it is useful to present the data in Fig. 14 by plotting $1/Z$ on a logarithmic scale, as shown in Fig. 15. Note that on this scale the exponential suppression of Z at strong coupling corresponds to a straight line. Obviously, for




FIG. 15: (Color online) Redrawing of Fig. 14: the inverse wave function renormalization factor $1/Z$ is now plotted on a logarithmic scale.




FIG. 16: (Color online) Low energy behavior of the spectral density $A(\omega)$ of the d -electrons as defined in Eqs. (4.42) for $u_0 = 0$ (dashed line) and $2, 4, 10$.

$U \gtrsim 15\Delta$ our two-channel FRG results begin to deviate significantly from the NRG results and definitely do not reproduce the known exponential suppression of the quasiparticle weight for $U \rightarrow \infty$. On the other hand, the two-channel FRG is accurate up to $U \approx 15\Delta$. We suspect that the deviations from the NRG results for $U \lesssim 15\Delta$ are due to our linearization of the spin susceptibility, which is expected to lead to a suppression of the quasiparticle weight, see Fig. 4.

Finally, we show in Fig. 16 the spectral density of the d -electrons, which is defined by

$$A(\omega) = -\frac{1}{\pi} \text{Im} G(\omega + i0). \quad (4.41)$$

For simplicity, we approximate $G(\omega + i0)$ by $G_{\Lambda=\omega}(i\omega \rightarrow \omega + i0)$ and obtain

$$\pi\Delta A(\omega) = \frac{1}{1 + \left(\frac{\omega}{Z(\omega)\Delta}\right)^2}, \quad (4.42)$$

with $Z(\omega) = Z_{l=\ln(\Lambda_0/\omega)}$. The qualitative behavior of

$A(\omega)$ is in agreement with known results¹: the width of the central Kondo peak is proportional to the wave function renormalization factor Z and its height at zero frequency is pinned to the value $1/(\pi\Delta)$ in the particle-hole symmetric case considered here, so that the overall spectral weight of the low-energy peak is of order Z . The spectral line-shape at energies larger than Δ (including the broadened Hubbard bands at energy scales $\pm U/2$) cannot be described within our low-energy truncation.

V. SUMMARY AND CONCLUSIONS

In summary, we have proposed a functional integral approach to the Anderson impurity model which is based on the mapping of the original fermionic problem onto a mixed Bose-Fermi theory where the spin fluctuations are represented by bosonic Hubbard-Stratonovich fields. Our approach can be used to approximately calculate the spectral properties including the quasiparticle weight up to couplings $U \lesssim 15\Delta$.

This work also contains several technical advances which will be useful for other FRG calculations. In particular, we have shown that in FRG calculations for Fermi systems using the technique of partial bosonization,^{17,18,19,30,31,32} the skeleton equations in the bosonic sector (which follow from the Dyson-Schwinger equations) can be used to close the FRG flow equations in the fermionic sector. Moreover, we have shown how the ambiguities inherent in multi-field Hubbard-Stratonovich decouplings can be resolved if one demands minimal sensitivity with respect to small variations of a given choice of decoupling. For a symmetric Anderson impurity model, this naturally leads to the manifestly spin-rotationally symmetric decoupling (4.3) at strong coupling. With our decoupling scheme, this symmetry is

also respected by the truncated flow equations.

Our approach still has two major shortcomings:

- (i) We have not been able to reproduce the exponential dependence $Z \propto \exp[-\pi^2 u_0/8]$ of the wave function renormalization factor for $U \rightarrow \infty$. Although we have tried several modifications of our approach, the strong fluctuations responsible for an exponential suppression of Z for $U \rightarrow \infty$ are apparently not correctly described within our truncation of the exact FRG flow equations.
- (ii) In order to make progress without using complicated numerical methods, we have made approximations which are only accurate at low energies $|\omega| \lesssim \Delta$. However, a direct numerical solution of our FRG flow equations should give rise to a much better agreement with NRG data at small couplings U and might also correctly reproduce the high-energy behavior of the spectral function. Therefore one should numerically solve the coupled system of integro-differential equations given by Eqs. (3.12), (3.13) or Eqs. (4.19)–(4.23), without approximating the fermionic Green function on the right-hand side of these equations by the Fermi liquid form (3.16). The fact that with our simple approximations we get results which agree reasonably well with NRG results gives us hope that solving this coupled system of integro-differential equations could lead to very accurate results.

ACKNOWLEDGMENTS

We would like to acknowledge useful conversations with W. Hofstetter, T. Pruschke and A. Sinner. Part of the work by P. K. was performed during a visit at the International Center for Condensed Matter Physics at the University of Brasília which was financially supported by the DAAD/PROBRAL program. J. J. R. C. gratefully acknowledges a DAAD postdoc fellowship.

-
- ¹ A. C. Hewson, *The Kondo Problem to Heavy Fermions*, (Cambridge University Press, Cambridge, 1993).
 - ² P. W. Anderson, Phys. Rev. B **124**, 41 (1961); Rev. Mod. Phys. **50**, 191 (1978).
 - ³ K. G. Wilson, Rev. Mod. Phys. **47**, 773 (1975).
 - ⁴ T. A. Costi, A. C. Hewson, and V. Zlatić, J. Phys.: Condens. Matter **6**, 2519 (1994).
 - ⁵ W. Hofstetter, Phys. Rev. Lett. **85**, 1508 (2000).
 - ⁶ R. Bulla, T. A. Costi, and T. Pruschke, Rev. Mod. Phys. **80**, 395 (2008).
 - ⁷ A. Georges, G. Kotliar, W. Krauth, and M. J. Rozenberg, Rev. Mod. Phys. **68**, 13 (1996).
 - ⁸ D. M. Newns and J. Read, J. Phys. C **16**, 3273 (1983); Adv. Phys. **36**, 799 (1988).
 - ⁹ P. Coleman, Phys. Rev. B **29**, 2 (1984).
 - ¹⁰ E. Mueller-Hartmann, Z. Phys. B **57**, 281 (1984).
 - ¹¹ D. E. Logan, M. P. Eastwood, and M. A. Tusch, J. Phys.: Condens. Matter **10**, 2673 (1998); N. L. Dickens and D. Logan, *ibid.* **13**, 4505 (2001).
 - ¹² A. C. Hewson, J. Phys.: Condens. Matter **13**, 10011 (2001).
 - ¹³ J. Kroha and P. Wölfle, in *Theoretical Methods for Strongly Correlated Electrons*, edited by D. Sénéchal, A.-M. Tremblay, and C. Bourbonnais (Springer, Berlin, 2004).
 - ¹⁴ V. Janiš and P. Augustinský, Phys. Rev. B **75**, 165108 (2007); *ibid.* **77**, 85106 (2008).
 - ¹⁵ R. Hedden, V. Meden, T. Pruschke, and K. Schönhammer, J. Phys.: Condens. Matter **16**, 5279 (2004).
 - ¹⁶ C. Karrasch, R. Hedden, R. Peters, T. Pruschke, K. Schönhammer, and V. Meden, J. Phys.: Condens. Matter **20**, 345205 (2008).
 - ¹⁷ F. Schütz, L. Bartosch, and P. Kopietz, Phys. Rev. B **72**, 035107 (2005).
 - ¹⁸ F. Schütz and P. Kopietz, J. Phys. A: Math. Gen. **39**, 8205 (2006).
 - ¹⁹ P. Kopietz, L. Bartosch, and F. Schütz, *Lectures on the Renormalization Group — from the Foundations to the Functional Renormalization Group*, (Springer, Berlin, 2009).
 - ²⁰ S. Ledowski and P. Kopietz, Phys. Rev. B **75**, 045134 (2007).

- (2007).
- ²¹ D. Litim, Phys. Rev. D **64**, 105007 (2001).
 - ²² T. R. Morris, Int. J. Mod. Phys. A **9**, 2411 (1994).
 - ²³ This result is implicitly contained in A. M. Tsvelick and P. B. Wiegmann, Adv. Phys. **32**, 453 (1983).
 - ²⁴ D. R. Hamann, Phys. Rev. Lett. **23**, 95 (1969).
 - ²⁵ S. Q. Wang, W. E. Evenson, and J. R. Schrieffer, Phys. Rev. Lett. **23**, 92 (1969).
 - ²⁶ C. Castellani and C. Di Castro, Phys. Lett. **70A**, 37 (1979).
 - ²⁷ H. J. Schulz, Phys. Rev. Lett. **65**, 2462 (1990).
 - ²⁸ C. A. Macêdo and M. D. Coutinho-Filho, Phys. Rev. B **43**, 13515 (1991).
 - ²⁹ N. Dupuis, Phys. Rev. B **65**, 245118 (2002); K. Borejsza and N. Dupuis, Europhys. Lett. **63**, 722 (2003); N. Dupuis, Phys. Rev. A **72**, 013606 (2005).
 - ³⁰ T. Baier, E. Bick, and C. Wetterich, Phys. Rev. B **70**, 125111 (2004); Phys. Lett. B **605**, 144 (2005).
 - ³¹ C. Wetterich, Phys. Rev. B **75**, 85102 (2007).
 - ³² P. Strack, R. Gersch, and W. Metzner, Phys. Rev. B **78**, 014522 (2008).